

Review

Can chromosomal instability initiate tumorigenesis?

Franziska Michor^a, Yoh Iwasa^b, Bert Vogelstein^c, Christoph Lengauer^c, Martin A. Nowak^{a,*}

^a Program for Evolutionary Dynamics, Department of Organismic and Evolutionary Biology, Department of Mathematics, Harvard University, Cambridge, MA 02138, USA

^b Department of Biology, Kyushu University, Fukuoka 812-8581, Japan

^c Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University, Baltimore, MD 21231, USA

Abstract

Cancers result from the accumulation of inherited and somatic mutations in oncogenes and tumor suppressor genes. These genes encode proteins that function in growth regulatory and differentiation pathways. Mutations in those genes increase the net reproductive rate of cells. Chromosomal instability (CIN) is a feature of most human cancers. Mutations in CIN genes increase the rate at which whole chromosomes or large parts of chromosomes are lost or gained during cell division. CIN causes an imbalance in chromosome number (aneuploidy) and an enhanced rate of loss of heterozygosity, which is an important mechanism of inactivating tumor suppressor genes. A crucial question of cancer biology is whether CIN is an early event and thus a driving force of tumorigenesis. Here we discuss mathematical models of situations where inactivation of one or two tumor suppressor genes is required for tumorigenesis. If two tumor suppressor genes have to be inactivated in rate-limiting steps, then CIN is likely to emerge before the inactivation of the first tumor suppressor gene.

© 2004 Elsevier Ltd. All rights reserved.

Keywords: Somatic evolution of cancer; Mathematical model; Chromosomal instability; Tumor suppressor gene inactivation

Contents

1. Introduction	43
2. Chromosomal instability before one tumor suppressor gene	44
3. Chromosomal instability before two tumor suppressor genes	45
4. Conclusions	47
5. Methods	47
Acknowledgements	48
References	48

1. Introduction

Tumor suppressor genes (TSGs) are negative regulators of cellular growth [1–5]. Loss of their function contributes to tumorigenesis [6–8]. The concept of a TSG emerged from a statistical analysis of retinoblastoma incidence in children [6]. This study and subsequent work led to the 2-hit hypothesis, which proposes that two hits in the retinoblastoma gene

are the rate limiting steps of this particular cancer [5]. In the inherited form, the first mutation is present in the germ line, whereas the second mutation emerges during somatic cell divisions. In the sporadic form, both mutations arise during somatic cell divisions. Inactivation of the first allele of a TSG does not normally change the phenotype of the cell and is considered an (almost) neutral mutation. Inactivation of the second allele confers a fitness advantage to the cell and can promote neoplastic growth. Human cancers result from the accumulation of mutations in several TSGs, oncogenes, and genes that are involved in maintaining genomic stabil-

* Corresponding author.

E-mail address: martin.nowak@harvard.edu (M.A. Nowak).

ity. Here, we discuss mathematical models of cancers that require the inactivation of one or two TSGs in rate limiting steps.

Genetic instability is a defining characteristic of most human cancers [9,10]. Two types of genetic instability have been identified [11]. In a small fraction of colorectal and some other cancers, a defect in mismatch repair results in an elevated mutation rate at the nucleotide level and consequent widespread microsatellite instability (MIN) [12,13]. The majority of colorectal and most other solid cancers, however, have chromosomal instability (CIN). CIN refers to an increased rate of losing or gaining whole chromosomes or large parts of chromosomes during cell division. The consequence of CIN is an imbalance in chromosome number (aneuploidy) and an increased rate of loss of heterozygosity (LOH). An elevated rate of LOH is an important property of CIN, because it accelerates the inactivation of TSGs.

A large number of genes that trigger CIN when mutated have been identified in the yeast *Saccharomyces cerevisiae* [14–16]. These so-called ‘CIN genes’ are involved in chromosome condensation, sister-chromatid cohesion, kinetochore structure and function, and microtubule formation as well as in cell cycle checkpoints. By comparison with yeast, we expect several hundred human CIN genes, but only few have been identified so far [17]. These genes include hBUB1, MAD2, BRCA1, BRCA2 and hCDC4 [18–22]. The genes hBUB1 and MAD2 encode proteins that are required for function of the spindle assembly checkpoint [18,20]. This checkpoint modulates the timing of anaphase initiation in mitotic cells containing improperly aligned chromosomes and increases the probability of successful delivery of a correct chromosome set to each daughter cell. The genes BRCA1 and BRCA2 encode proteins that are implicated in DNA repair and recombination, checkpoint control of the cell cycle, and transcription [19,21]. hCDC4 is an E3 ubiquitin ligase that is thought to be involved in regulating the G1-S cell-cycle checkpoint by targeting proteins for destruction [22].

The classification of CIN genes is based on the mutational events required to engage CIN [10]. Class I CIN genes, such as MAD2, trigger CIN if one allele of the gene is mutated or lost. Class II CIN genes, such as hBUB1, trigger CIN if one allele is mutated in a dominant negative fashion. Both class I and class II. CIN genes are ‘single hit’ CIN genes. Class III CIN genes, such as BRCA1 and BRCA2, trigger CIN if both alleles are mutated.

The mathematical investigation of cancer began in the 1960s, when Nordling [23], Armitage and Doll [24,25], and Fisher [26] set out to explain the age-dependent incidence curves of human cancers. These seminal studies led to the idea that multiple probabilistic events are required for the somatic evolution of cancer [27,28]. In the early 1980s, Knudson used a statistical analysis of the incidence of retinoblastoma in children to explain the role of tumor suppressor genes in sporadic and inherited cancers [6]. This work was later extended to a two-stage stochastic model for the process of cancer initiation and progression [29], which inspired much subsequent




Fig. 1. Inactivation dynamics of a tumor suppressor gene (TSG), A , with and without chromosomal instability (CIN). (a) The wild type cell ('start') has two unmutated alleles of TSG A , $A^{+/+}$. The cell evolves from $A^{+/+}$ via $A^{+/-}$ to $A^{-/-}$. CIN can arise at any stage of TSG inactivation and causes very fast LOH. (b) A compartment of cells evolves along the evolutionary trajectory. If an $A^{+/-}$ cell clone with CIN produces an $A^{-/-}$ cell before taking over the compartment, a stochastic tunnel arises (diagonal arrow); the compartment evolves from $A^{+/-}$ without CIN to $A^{-/-}$ with CIN without ever visiting $A^{+/-}$ with CIN.

work [30–32]. Later on, specific theories for drug resistance [33,34], angiogenesis [35], immune responses against tumors [36], and genetic instabilities [37–40] were developed.

A major question in cancer genetics is to what extent CIN, or any genetic instability, is an early event and thus a driving force in tumorigenesis [41–47]. Here, we discuss mathematical models that investigate the role of CIN in cancers which require elimination of one or two TSGs [42,46–48].

2. Chromosomal instability before one tumor suppressor gene

Consider a path to cancer where both alleles of a TSG, A , have to be inactivated in a single cell (Fig. 1a) [42,46–48]. Initially, the cell is wild type, $A^{+/+}$. The cell evolves from $A^{+/+}$ via $A^{+/-}$ to $A^{-/-}$. The first allele of a TSG is typically inactivated by a point mutation. The second allele can be inactivated by one of several possible mechanisms (Fig. 2). It can be inactivated by a second point mutation, leading to two distinct point mutations in the two alleles. Alternatively, the chromosome on which the second allele resides can be lost; this chromosome loss might be deleterious or lethal. The two alleles at the TSG locus can recombine mitotically, leading to two identical point mutations on partly homozygous alleles. Finally, the chromosome harboring the wild type allele can be lost and the chromosome harboring the first point mutation duplicated. This mechanism may require two steps, occurring in two distinct cell divisions, but could also occur in a single division through a process broadly related to mitotic recombination.




Fig. 2. Mechanisms of tumor suppressor gene (TSG) inactivation. The first TSG allele is typically inactivated by a point mutation. The second allele can be inactivated by a second point mutation, by chromosome loss, mitotic recombination, or chromosome loss followed by duplication of the mutated allele.

CIN can emerge at any stage of tumorigenesis. First, assume that CIN is dominant, i.e. triggered by mutation of a class I or II CIN gene [10,42]. The crucial effect of CIN is to increase the rate of LOH, thereby accelerating the transition from $A^{+/-}$ to $A^{-/-}$. CIN can have a cost for the cell by increasing the rate of accumulating lethal mutations and triggering apoptosis. CIN, however, can also be advantageous by reducing the duration of the cell cycle (if certain checkpoints have been eliminated) or by providing increased evolvability in detrimental environments [49,50]. Therefore, it makes sense to consider possibilities of costly, neutral, and advantageous CIN phenotypes.

Tissues are organized into small compartments of cells [51–54]. Not all cells of a compartment might be at risk of becoming cancer cells, however, because certain mutations might only have an effect if they occur in stem cells [55]. Suppose N_0 cells are at risk in any one compartment. If the mutation rate is smaller than the inverse of N_0 , then the approximation of homogeneous compartments holds: a mutated cell will either take over the compartment or go extinct before the next mutation arises. If an $A^{+/-}$ cell clone with CIN produces an $A^{-/-}$ cell before taking over the compartment, then the phenomenon of ‘stochastic tunneling’ arises [42,56]: the compartment moves from $A^{+/-}$ without CIN to $A^{-/-}$ with CIN without ever visiting $A^{+/-}$ with CIN. If CIN is neutral or advantageous, then the tunnel does not arise. Mathematical procedures are outlined in Section 5.

3. Chromosomal instability before two tumor suppressor genes

Consider a path to cancer where two TSGs, A and B , have to be inactivated in a single cell (Fig. 3a) [42,46–48]. Initially, the cell is wild type, $A^{+/+}B^{+/+}$. Suppose gene

A has to be inactivated first. Hence the cell evolves from $A^{+/+}B^{+/+}$ via $A^{+/-}B^{+/+}$ to $A^{-/-}B^{+/+}$, and subsequently to $A^{-/-}B^{+/-}$ and $A^{-/-}B^{-/-}$. CIN can emerge at any stage of tumorigenesis due to mutations of class I, II or III CIN genes. Once CIN has emerged, it accelerates the transitions from $A^{+/-}$ to $A^{-/-}$ and from $B^{+/-}$ to $B^{-/-}$.

Evolutionary dynamics within a compartment of N_0 cells are illustrated in Fig. 3b. CIN might emerge before gene A has been inactivated. A stochastic tunnel arises if an $A^{+/-}B^{+/+}$ cell clone with CIN produces an $A^{-/-}B^{+/+}$ cell before taking over the compartment [42,56]. Then the compartment moves from $A^{+/-}B^{+/+}$ without CIN to $A^{-/-}B^{+/+}$ with CIN without ever visiting $A^{+/-}B^{+/+}$ with CIN.

Inactivation of the first TSG can induce neoplastic growth. We assume that the $A^{-/-}$ compartment gives rise to a small lesion of N_1 cells. In this lesion the second TSG has to be inactivated for further tumor progression. Due to the increased compartment size, the evolutionary pathway might tunnel from $A^{-/-}B^{+/+}$ directly to $A^{-/-}B^{-/-}$. This means that the $A^{-/-}B^{+/-}$ cell does not reach fixation before the $A^{-/-}B^{-/-}$ cell arises. Mathematical procedures are outlined in Section 5.

The importance of early CIN in tumorigenesis depends on the number of possible CIN mutations, which in turn depends on the number of CIN genes in the human genome. We can calculate the minimum number of CIN genes in the genome that are needed to ensure that a CIN mutation precedes the inactivation of one or two TSGs. Let us discuss some plausible parameter choices (Table 1). The mutation rate per base per cell division is 10^{-10} to 10^{-11} [57]. If an average TSG can be inactivated by any one of 1000 point mutations, the mutation rate per gene per cell division is $u = 10^{-7}$. Estimates of the rate of LOH in non-CIN cells range from $p_0 = 10^{-7}$ to $p_0 = 10^{-5}$. In our opinion, the most likely scenario is that p_0 has the same order of magnitude as u . In this case and in the absence of CIN, the hit inactivating the second TSG allele is sometimes LOH and sometimes a point mutation. If $p_0 \gg u$, then two distinct point mutations should never be observed in the two TSG alleles. The rate of LOH in CIN cells is $p = 10^{-2}$ [58]. Consider a compartment size of $N_0 = 4$. This choice is motivated by the geometry of the colonic crypt [54]. Suppose the size of the lesion after clonal expansion of an $A^{-/-}$ cell clone is of the order of $N_1 = 10^4$ actively dividing cells. CIN can be disadvantageous, neutral, or advantageous [17,49]. The most substantial cost CIN can possibly have is $r = (1 - p)^{45} \approx 0.6$; this means that loss of any chromosome other than the one containing the TSG locus is lethal. CIN is neutral if $r = 1$ and advantageous if $r > 1$. Given these parameter choices and $r = 0.6$, we find that 3 class I CIN genes, or 29 class II CIN genes, or more than 100 class III CIN genes are required to ensure that CIN precedes inactivation of the first TSG in a pathway where no other TSG has to be eliminated in a rate limiting step. If two TSGs have to be inactivated in rate limiting steps, then 1 class I CIN gene, or 1 class II CIN gene, or 30 class III CIN genes are sufficient for CIN to precede inactivation of the first




Fig. 3. Mutational network of inactivating two tumor suppressor genes (TSGs), A and B, with and without chromosomal instability (CIN). (a) The wild type cell ('start') has two unmutated alleles of both TSGs, $A^{+/+} B^{+/+}$. Mutations lead via $A^{+/-} B^{+/+}$, $A^{-/-} B^{+/+}$ and $A^{-/-} B^{+/-}$ to $A^{-/-} B^{-/-}$. Without CIN, the cell maintains a stable genotype (second row). CIN can arise at any stage of tumorigenesis due to mutations in a class I, II, or III CIN gene. Classes I and II CIN genes require one hit to trigger CIN (first row). Class III CIN genes require two hits to trigger CIN, one in each allele (third and fourth row). Once arisen, CIN accelerates the inactivation of the second allele of each following TSG. CIN, however, has a cost for the cell by increasing the chance of lethal mutations and apoptosis. (b) Inactivation dynamics starting with a population of N_0 wild type cells. CIN can emerge at any stage of tumorigenesis. If an $A^{+/-} B^{+/+}$ cell clone without CIN produces an $A^{-/-} B^{+/+}$ cell with CIN before reaching fixation, the compartment tunnels to $A^{-/-} B^{+/+}$ with CIN (red arrows). Inactivation of TSG A leads to a small lesion of N_1 cells in which TSG B is eliminated. Due to the increased compartment size, the evolutionary pathway tunnels directly to $A^{-/-} B^{-/-}$ (red arrows).

TSG. Table 1 provides further examples for other parameter choices.

The parameters have different effects on the importance of the early CIN. An increase in N_0 , N_1 and p_0 as well as a decrease in r and p make the early emergence of CIN less likely. Instead of calculating the inactivation dynamics of one specific TSG, we can also study the possibility that one gene out of a family of TSGs can be inactivated. Suppose gene A

is any one of ten TSGs and likewise gene B is any one of ten TSGs. This is an extreme example. In this scenario and using the same parameter values as above, the same number of CIN genes in the human genome as before is required to ensure that CIN precedes inactivation of the first TSG in a pathway where no other TSG has to be eliminated in a rate limiting step. If two TSGs have to be inactivated in rate limiting steps, then 1 class I CIN gene, or 5 class II CIN gene, or 62 class

Table 1
Minimum number of CIN genes needed to ensure that CIN emerges before one or two TSGs

		CIN before 1 TSG			CIN before 2 TSGs						
		independent of N_1			$N_1 = 10^4$			$N_1 = 10^5$			
		r	n_1	n_2	n_3	n_1	n_2	n_3	n_1	n_2	n_3
(a)	1.4	1	2	95	1	1	8	1	1	21	
	1.0	2	3	>100	1	1	11	1	1	26	
	0.6	3	6	>100	1	1	20	1	1	42	
(b)	1.4	1	2	>100	1	1	10	1	1	23	
	1.0	3	6	>100	1	1	17	1	1	40	
	0.6	27	54	>100	1	1	59	2	4	>100	
(c)	1.4	1	8	>100	1	1	13	1	2	31	
	1.0	2	12	>100	1	1	16	1	2	38	
	0.6	3	29	>100	1	1	30	1	5	61	
(d)	1.4	1	66	>100	1	5	24	1	27	65	
	1.0	2	>100	>100	1	8	31	1	40	76	
	0.6	3	>100	>100	1	25	54	1	100	>100	

Class I CIN genes, n_1 , trigger CIN if one allele is mutated or lost. Class II CIN genes, n_2 , trigger CIN if one allele is mutated in a dominant negative fashion. Class III CIN genes, n_3 , trigger CIN if both alleles are mutated. The somatic fitness of CIN cells is denoted by r , and the compartment size after clonal expansion by N_1 . Results are obtained by numerical simulation of Eq. (1). Parameter values are $u = 10^{-7}$, $p = 10^{-2}$, $t = 80$ years, and $p_0 = 10^{-7}$ and $N_0 = 4$ in (a); $p_0 = 10^{-7}$ and $N_0 = 10$ in (b); $p_0 = 10^{-6}$ and $N_0 = 4$ in (c); and $p_0 = 10^{-5}$ and $N_0 = 4$ in (d).

III CIN genes are sufficient for CIN to precede inactivation of the first TSG. Hence, a large number of alternative TSGs increases the number of classes II and III CIN genes needed, but does not significantly alter the number of class I CIN genes required for CIN to arise early.

4. Conclusions

Even if many alternative CIN genes are needed to ensure that costly CIN emerges before one TSG, only very few CIN genes suffice for CIN to precede two TSGs. This effect is especially strong if inactivation of the first TSG leads to a moderate clonal expansion. In this case, the second TSG must be inactivated in a rate limiting fashion. If, on the other hand, the inactivation of the first TSG causes a rapid clonal expansion ($N_1 \gg 10^5$), then the inactivation of the second TSG does not occur in a rate limiting step, and CIN can only accelerate inactivation of the first TSG. A wide range of plausible parameter values, which are conservatively biased against CIN, all give the same message: of the order of 1 (to 10) neutral CIN genes are needed to ensure that CIN emerges before the inactivation of one TSG; of the order of 1 (to 10) costly CIN genes are needed to ensure that CIN emerges before the inactivation of the first of two TSGs. By analogy with yeast, we expect several hundreds of CIN genes in the human genome. Therefore it is likely that in any one human tissue a large number of genes will lead to CIN

when mutated. Thus, in any pathway of cancer progression where at least two TSGs need to be eliminated in rate limiting steps, CIN will arise before inactivation of the first TSG and therefore initiate the mutational sequence that leads to cancer.

5. Methods

The stochastic process illustrated in Figs. 1b and 3b can be described by differential equations. Denote by X_0, X_1, X_2 , and X_4 the probabilities that a compartment is in state $A^{+/+}B^{+/+}$, $A^{+/-}B^{+/+}$, $A^{-/-}B^{+/+}$, and $A^{-/-}B^{-/-}$, respectively, without chromosomal instability (CIN). Denote by Y_0, Y_1, Y_2 , and Y_4 the probabilities that a compartment is in state $A^{+/+}B^{+/+}$, $A^{+/-}B^{+/+}$, $A^{-/-}B^{+/+}$, and $A^{-/-}B^{-/-}$, respectively, with CIN. The differential equations are given by

$$\begin{aligned}
 \dot{X}_0 &= X_0(-r_0 - c_0) & \dot{Y}_0 &= c_0X_0 - s_0Y_0 \\
 \dot{X}_1 &= r_0X_0 - X_1(r_1 + c_0 + c_1) & \dot{Y}_1 &= s_0Y_0 + c_0X_1 - s_1Y_1 \\
 \dot{X}_2 &= r_1X_1 - r_2X_2 & \dot{Y}_2 &= s_1Y_1 + c_1X_1 - s_2Y_2 \\
 \dot{X}_4 &= r_2X_2 & \dot{Y}_4 &= s_2Y_2
 \end{aligned}
 \tag{1}$$

The transition rates between the states are given by: $r_0 = 2u/\tau_0$, $r_1 = N_0(u + p_0)/\tau_0$, $r_2 = N_12u\sqrt{u + p_0}/\tau_1$, $c_0 = N_0u_c\rho/\tau_0$, $c_1 = [N_0u_c r/(1 - r)][p/\tau_0]$, $s_0 = r_0$, $s_1 = N_0p/\tau_0$, and $s_2 = rN_12u\sqrt{p}/\tau_1$ [42,46–48]. Cells with genotype $A^{+/+}B^{+/+}$ and $A^{+/-}B^{+/+}$ divide every τ_0 days, while cells with genotype $A^{-/-}B^{+/+}$ and $A^{-/-}B^{+/-}$ divide every τ_1 days. Let us first discuss the mutation–selection network leading from X_0 via X_1 and X_2 to X_4 . The rate of change, r_0 , is equal to the rate that a mutation occurs in the first allele of TSG A times the probability that the mutant cell will take over the compartment. The mutation occurs at rate $2N_0u/\tau_0$, because (i) the mutation can arise in any one of N_0 cells; (ii) the cells divide once every τ_0 days; (iii) the mutation rate per gene per cell division is given by u ; and (iv) there are two alleles. The probability that the mutated cell takes over the compartment (i.e. reaches fixation) is given by $1/N_0$, if we assume that the mutation is neutral [59]. Thus, the population size, N_0 , cancels in the product, and we simply obtain $2u/\tau_0$. Once the first allele of TSG A has been inactivated, the second allele can be inactivated either by another point mutation or an LOH event. This process occurs at rate $r_1 = N_0(u + p_0)/\tau_0$, where p_0 is the rate of LOH in normal cells. We assume that $A^{-/-}$ cells have a large fitness advantage, which means the probability of fixation of such a cell is close to one. Inactivation of both alleles of TSG A leads to a small lesion of N_1 cells dividing every τ_1 days. In this lesion, TSG B is inactivated at rate $r_2 = N_12u\sqrt{u + p_0}/\tau_1$ [42,46–48]. CIN can arise at any stage of tumorigenesis and causes very fast LOH [58]. Denote the rate of LOH in CIN

cells by p ; we have $p \gg p_0$. The rate of change, c_0 , is equal to the rate at which a mutation triggering CIN occurs, $N_0 u_c \tau_0$, times the probability that the mutant cell will take over the compartment. The mutation rate triggering CIN, u_c , depends on the number of available classes I, II and III CIN genes in the human genome, n_1 , n_2 and n_3 , respectively. The mutation rates of classes I and II CIN genes are $u_c = 2n_1(u + p_0)$ and $u_c = 2n_2u$, respectively. The mutation rate of the first allele of class III CIN genes is $2n_3u$, and the mutation rate of the second allele of class III CIN genes is $u + p_0$. The probability of fixation of a CIN cell depends on its somatic fitness, r , and is given by $\rho = (1 - 1/r)/(1 - 1/r^{N_0})$. This is the standard fixation probability of a Moran process [60]. Therefore, the rate of change is given by $c_0 = N_0 u_c \rho / \tau_0$. A stochastic tunnel arises if an $A^{-/-} B^{+/+}$ cell with CIN reaches fixation before an $A^{+/-} B^{+/+}$ cell does. The rate of tunneling is given by $c_1 = [N_0 u_c r / (1 - r)] [p / \tau_0]$.

Let us now discuss the mutation–selection network leading from Y_0 via Y_1 and Y_2 to Y_4 . The rate of change, s_0 , is equal to the rate r_0 , because CIN does not change the probability of point mutations. The rate at which the second allele of TSG A is inactivated is given by $s_1 = N_0 p / \tau_0$. The rate at which TSG B is inactivated is given by $s_2 = r N_1 2u \sqrt{p} / \tau_1$. Eq. (1) is a system of linear differential equations which can be solved analytically using standard techniques.

Acknowledgements

The Program for Evolutionary Dynamics is supported by Jeffrey E. Epstein.

References

- [1] Friend SH, Bernards R, Rogelj S, Weinberg RA, Rapaport JM, Albert DM, et al. A human DNA segment with properties of the gene that predisposes to retinoblastoma and osteosarcoma. *Nature* 1986;323:643–6.
- [2] Kinzler KW, et al. Identification of a gene located at chromosome 5q21 that is mutated in colorectal cancers. *Science* 1991;251:1366–70.
- [3] Weinberg RA. Tumor suppressor genes. *Science* 1991;254:1138–46.
- [4] Knudson AG. Antioncogenes and human cancer. *Proc Natl Acad Sci USA* 1993;90:10914–21.
- [5] Vogelstein B, Kinzler KW. The genetic basis of human cancer. 2nd ed. Toronto: McGraw-Hill; 2001.
- [6] Knudson AG. Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci USA* 1971;68:820–3.
- [7] Luebeck EG, Moolgavkar SH. Multistage carcinogenesis and the incidence of colorectal cancer. *Proc Natl Acad Sci USA* 2002;99:15095–100.
- [8] Park BH, Vogelstein B. In: Abeloff MD, editor. *Clinical oncology*. 2nd ed. Churchill Livingstone; 2000.
- [9] Loeb LA. A mutator phenotype in cancer. *Cancer Res* 2001;61:3230–9.
- [10] Rajagopalan H, Nowak MA, Vogelstein B, Lengauer C. The significance of unstable chromosomes in colorectal cancer. *Nat Rev Can* 2003;3:695–701.
- [11] Lengauer C, Kinzler KW, Vogelstein B. Genetic instabilities in human cancers. *Nature* 1998;396:623–49.
- [12] Kinzler KW, Vogelstein B. Lessons from hereditary colon cancer. *Cell* 1996;87:159–70.
- [13] Perucho M. Cancer of the microsatellite phenotype. *Biol Chem* 1996;377:675–84.
- [14] Shonn MA, McCarroll R, Murray AW. Requirement of the spindle checkpoint for proper chromosome segregation in budding yeast meiosis. *Science* 2000;289:300–3.
- [15] Kolodner RD, Putnam CD, Myung K. Maintenance of genome stability in *Saccharomyces cerevisiae*. *Science* 2002;297:552–7.
- [16] Nasmyth K. Segregating sister genomes: the molecular biology of chromosome separation. *Science* 2002;297:559–65.
- [17] Cahill DP, Lengauer C, Yu J, Riggins GJ, Willson JKV, Markowitz SD, et al. Mutations of mitotic checkpoint genes in human cancers. *Nature* 1998;392:300–3.
- [18] Li Y, Benzra R. Identification of a human mitotic checkpoint gene: *hsMAD2*. *Science* 1996;274:246–8.
- [19] Milner J, Ponder B, Hughes-Davies L, Seltnann M, Kouzarides T. Transcriptional activation functions in *BRCA2*. *Nature* 1997;386:772–3.
- [20] Pangilinan F, Li Q, Weaver T, Lewis BC, Dang CV, Spencer F. Mammalian BUB1 protein kinases: map positions and in vivo expression. *Genomics* 1997;46:379–88.
- [21] Yarden RI, Pardo-Reoyo S, Sgagias M, Cowan KH, Brody LC. *BRCA1* regulates the G2/M checkpoint by activating Chk1 kinase upon DNA damage. *Nat Genet* 2002;30:265–9.
- [22] Rajagopalan H, Jallepalli PV, Rago C, Velculescu VE, Kinzler KW, Vogelstein B, et al. Inactivation of *hCDC4* can cause chromosomal instability. *Nature* 2004;428:77–81.
- [23] Nordling CO. A new theory on cancer-inducing mechanism. *Br J Cancer* 1953;7:68–72.
- [24] Armitage P, Doll R. The age distribution of cancer and a multi-stage theory of carcinogenesis. *Br J Cancer* 1954;8:1–12.
- [25] Armitage P, Doll R. A two-stage theory of carcinogenesis in relation to the age distribution of human cancer. *Br J Cancer* 1957;11:161–9.
- [26] Fisher JC. Multiple-mutation theory of carcinogenesis. *Nature* 1959;181:651–2.
- [27] Nunney L. Lineage selection and the evolution of multistage carcinogenesis. *Proc R Soc Lond B* 1999;266:493–8.
- [28] Tomlinson I, Sasienski P, Bodmer W. How many mutations in a cancer? *Am J Pathol* 2002;160:755–8.
- [29] Moolgavkar SH, Knudson AG. Mutation and cancer: a model for human carcinogenesis. *J Natl Cancer Inst* 1981;66:1037–52.
- [30] Luebeck EG, Moolgavkar SH. Multistage carcinogenesis and the incidence of colorectal cancer. *Proc Natl Acad Sci USA* 2002;99:15095–100.
- [31] Grist SA, McCarron M, Kutlaca A, Turner DR, Morley AA. In vivo human somatic mutation: frequency and spectrum with age. *Mutat Res* 1992;266:189–96.
- [32] Gatenby RA, Vincent TL. An evolutionary model of carcinogenesis. *Cancer Res* 2003;63:6212–20.
- [33] Goldie JH, Coldman AJ. A mathematic model for relating the drug sensitivity of tumors to their spontaneous mutation rate. *Cancer Treat Rep* 1979;63:1727–33.
- [34] Goldie JH, Coldman AJ. Quantitative model for multiple levels of drug resistance in clinical tumors. *Cancer Treat Rep* 1983;67:923–31.
- [35] Anderson AR, Chaplain MA. Continuous and discrete mathematical models of tumor-induced angiogenesis. *Bull Math Biol* 1998;60:857–99.
- [36] Owen MR, Sherratt JA. Mathematical modeling of macrophage dynamics in tumors. *Math Models Methods Appl Biol Chem* 2004;377:675–84.
- [37] Taddei F, Radman M, Maynard-Smith J, Toupance B, Gouyon PH, Godelle B. Role of mutator alleles in adaptive evolution. *Nature* 1997;387:700–2.

- [38] Little MP, Wright EG. A stochastic carcinogenesis model incorporating genomic instability fitted to colon cancer data. *Math Biosci* 2003;183:111–34.
- [39] Breivik J, Gaudernack G. Resolving the evolutionary paradox of genetic instability: a cost–benefit analysis of DNA repair in changing environments. *FEBS Lett* 2004;563:7–12.
- [40] Sole RV, Deisboeck TS. An error catastrophe in cancer? *J Theor Biol* 2004;228:47–54.
- [41] Loeb LA. Mutator phenotype may be required for multistage carcinogenesis. *Cancer Res* 1991;51:3075–9.
- [42] Nowak MA, Komarova NL, Sengupta A, Jallepalli PV, Shih IeM, Vogelstein B, et al. The role of chromosomal instability in tumor initiation. *Proc Natl Acad Sci USA* 2002;99:16226–31.
- [43] Sieber OM, Heinemann K, Gorman P, Lamlum H, Crabtree M, Simpson CA, et al. Analysis of chromosomal instability in human colorectal adenomas with two mutational hits at APC. *Proc Natl Acad Sci USA* 2002;99:16910–5.
- [44] Little MP, Wright EG. A stochastic carcinogenesis model incorporating genomic instability fitted to colon cancer data. *Math Biosci* 2003;183:111–34.
- [45] Michor F, Iwasa Y, Komarova NL, Nowak MA. Local regulation of homeostasis favors chromosomal instability. *Curr Biol* 2003;13:581–4.
- [46] Komarova NL, Sengupta A, Nowak MA. Mutation–selection networks of cancer initiation: tumor suppressor genes and chromosomal instability. *J Theor Biol* 2003;223:433–50.
- [47] Michor F, Iwasa Y, Nowak MA. Dynamics of cancer progression. *Nat Rev Cancer* 2004;4:197–206.
- [48] Nowak MA, Michor F, Komarova NL, Iwasa Y. *Proc Natl Acad Sci USA* 2004;101:10635–8.
- [49] Breivik J. Don't stop for repairs in a war zone: Darwinian evolution unites genes and environment in cancer development. *Proc Natl Acad Sci USA* 2001;5379–81.
- [50] Bardelli A, Cahill DP, Lederer G, Speicher MR, Kinzler KW, Vogelstein B, et al. Carcinogen-specific induction of genetic instability. *Proc Natl Acad Sci USA* 2001;98:5770–5.
- [51] Mintz B. Clonal basis of mammalian differentiation. *Symp Soc Exp Biol* 1971;25:345–70.
- [52] Cairns J. Mutation selection and the natural history of cancer. *Nature* 1975;255:197–200.
- [53] Bach SP, Renehan AG, Potten CS. Stem cells: the intestinal stem cell as a paradigm. *Carcinogenesis* 2000;21:469–76.
- [54] Yatabe Y, Tavare S, Shibata D. Investigating stem cells in human colon by using methylation patterns. *Proc Natl Acad Sci USA* 2001;98:10839–44.
- [55] Michor F, Nowak MA, Frank SA, Iwasa Y. Stochastic elimination of cancer cells. *Proc R Soc Lond B* 2003;270:2017–24.
- [56] Iwasa Y, Michor F, Nowak MA. Stochastic tunnels in evolutionary dynamics. *Genetics* 2004;166:1571–9.
- [57] Kunkel TA, Bebenek K. DNA replication fidelity. *Annu Rev Biochem* 2000;69:497–529.
- [58] Lengauer C, Kinzler KW, Vogelstein B. Genetic instability in colorectal cancers. *Nature* 1997;386:623–7.
- [59] Kimura M. The role of compensatory neutral mutations in molecular evolution. *J Genet* 1985;64:7–19.
- [60] Nowak MA, Michor F, Iwasa Y. The linear process of somatic evolution. *Proc Natl Acad Sci USA* 2003;100:14966–9.